

# Moments of Traces for Circular $\beta$ -ensembles

Tiefeng Jiang

University of Minnesota

This is joint work with Sho Matsumoto

April 5, 2010

- Moments for Haar Unitary Matrices (D.E. Thm)
- Background for Circular  $\beta$ -Ensembles
- Moments for Circular  $\beta$ -Ensembles
- Proofs by Jack Polynomials

# 1. Moments for Haar Unitary Matrices

- ▶ What is Haar-invariant unitary matrix  $\Gamma_n$ ?

Mathematically,

$\Gamma_n$  : normalized Haar measure on  $U(n)$  : set of  $n$  by  $n$  unitary matrices.

# 1. Moments for Haar Unitary Matrices

## ► What is Haar-invariant unitary matrix $\Gamma_n$ ?

Mathematically,

$\Gamma_n$  : normalized Haar measure on  $U(n)$  : set of  $n$  by  $n$  unitary matrices.

Statistically,

Assume the entries of  $Y = Y_{n \times n}$  are i.i.d.  $\mathbb{C}N(0, 1)$ . Two ways to generate such matrices

# 1. Moments for Haar Unitary Matrices

► What is Haar-invariant unitary matrix  $\Gamma_n$ ?

Mathematically,

$\Gamma_n$  : normalized Haar measure on  $U(n)$  : set of  $n$  by  $n$  unitary matrices.

Statistically,

Assume the entries of  $Y = Y_{n \times n}$  are i.i.d.  $\mathbb{C}N(0, 1)$ . Two ways to generate such matrices.

# 1. Moments for Haar Unitary Matrices

## ► What is Haar-invariant unitary matrix $\Gamma_n$ ?

Mathematically,

$\Gamma_n$  : normalized Haar measure on  $U(n)$  : set of  $n$  by  $n$  unitary matrices.

Statistically,

Assume the entries of  $Y = Y_{n \times n}$  are i.i.d.  $\mathbb{C}N(0, 1)$ . Two ways to generate such matrices

1) The matrix  $Q$  in QR (Gram-Schmidt) decomposition of  $Y$

$$2) \Gamma_n \stackrel{d}{=} Y(Y^*Y)^{-1/2}$$

► Theorem (Diaconis and Evans: 2001)

(a)  $a = (a_1, \dots, a_k)$ ,  $b = (b_1, \dots, b_k)$  with  $a_j, b_j \in \{0, 1, 2, \dots, g\}$ .

$X_1, \dots, X_k$ : i.i.d.  $\mathbb{C}N(0, 1)$ . If  $n \sum_{j=1}^k |a_j - b_j| \rightarrow \infty$  as  $n \rightarrow \infty$ , then  $\mathbb{P}(X_1, \dots, X_k \in \text{support}(a)) \rightarrow \mathbb{P}(X_1, \dots, X_k \in \text{support}(b))$ .

► Theorem (Diaconis and Evans: 2001)

(a)  $a = (a_1, \dots, a_k)$ ,  $b = (b_1, \dots, b_k)$  with  $a_j, b_j \geq 0, 1, 2, \dots, g$ .  
 $X_1, \dots, X_k$ : i.i.d.  $\mathbb{C}N(0, 1)$ . If  $n \gg \sum_{j=1}^k ja_j - \sum_{j=1}^k jb_j$ ,

$$\mathbb{E} \left[ \prod_{j=1}^k (\text{Tr}(U_n^j))^{a_j} \overline{(\text{Tr}(U_n^j))^{b_j}} \right]$$

=



► Theorem (Diaconis and Evans: 2001)

(a)  $a = (a_1, \dots, a_k)$ ,  $b = (b_1, \dots, b_k)$  with  $a_j, b_j \geq 0, 1, 2, \dots, g$ .  
 $X_1, \dots, X_k$ : i.i.d.  $\mathbb{C}N(0, 1)$ . If  $n = \sum_{j=1}^k j a_j = \sum_{j=1}^k j b_j$ ,

$$\mathbb{E} \left[ \prod_{j=1}^k (\text{Tr}(U_n^j))^{a_j} \overline{(\text{Tr}(U_n^j))^{b_j}} \right]$$

$$= \delta_{ab} \prod_{j=1}^k j^{a_j} a_j!$$

► Theorem (Diaconis and Evans: 2001)

(a)  $a = (a_1, \dots, a_k)$ ,  $b = (b_1, \dots, b_k)$  with  $a_j, b_j \geq 0, 1, 2, \dots, g$ .  
 $X_1, \dots, X_k$ : i.i.d.  $\mathbb{C}N(0, 1)$ . If  $n = \sum_{j=1}^k j a_j = \sum_{j=1}^k j b_j$ ,

$$\begin{aligned} & \mathbb{E} \left[ \prod_{j=1}^k (\text{Tr}(U_n^j))^{a_j} \overline{(\text{Tr}(U_n^j))^{b_j}} \right] \\ &= \delta_{ab} \prod_{j=1}^k j^{a_j} a_j! = \delta_{ab} \mathbb{E} \left[ \prod_{j=1}^k (\sqrt{j} X_j)^{a_j} \overline{(\sqrt{j} X_j)^{b_j}} \right] \end{aligned}$$


► Theorem (Diaconis and Evans: 2001)

(a)  $a = (a_1, \dots, a_k)$ ,  $b = (b_1, \dots, b_k)$  with  $a_j, b_j \geq 0, 1, 2, \dots, g$ .  
 $X_1, \dots, X_k$ : i.i.d.  $\mathbb{C}N(0, 1)$ . If  $n = \sum_{j=1}^k j a_j = \sum_{j=1}^k j b_j$ ,


$$\begin{aligned} & \mathbb{E} \left[ \prod_{j=1}^k (\text{Tr}(U_n^j))^{a_j} \overline{(\text{Tr}(U_n^j))^{b_j}} \right] \\ &= \delta_{ab} \prod_{j=1}^k j^{a_j} a_j! = \delta_{ab} \mathbb{E} \left[ \prod_{j=1}^k (\sqrt{j} X_j)^{a_j} \overline{(\sqrt{j} X_j)^{b_j}} \right] \end{aligned}$$

(b) For  $j$  and  $k$ ,

$$\mathbb{E} [\text{Tr}(U_n^j) \overline{\text{Tr}(U_n^k)}] = \delta_{jk} j^{n/j}$$



*Circular Ensembles and Haar-invariant Matrices from Classical Compact Groups*



*Circular Ensembles and Haar-invariant Matrices from Classical Compact Groups*

Diaconis (2004) believes there is a good formula for  $COE$  and  $CSE$

## 2. Background for Circular $\beta$ -Ensembles

### ► Probability density function

$e^{i\theta_1}, \dots, e^{i\theta_n}$  : eigenvalues of Haar-invariant unitary matrix.

pdf:  $f(\theta_1, \dots, \theta_n | \beta = 2)$ , where

## 2. Background for Circular $\beta$ -Ensembles

### ► Probability density function

$e^{i\theta_1}, \dots, e^{i\theta_n}$  : eigenvalues of Haar-invariant unitary matrix.

pdf:  $f(\theta_1, \dots, \theta_n | \beta = 2)$ , where

$$f(\theta_1, \dots, \theta_n | \beta) = \text{Const} \prod_{1 \leq j < k \leq n} |j e^{i\theta_j} - k e^{i\theta_k}|^\beta$$

$\beta > 0, \theta_i \in [0, 2\pi)$

## 2. Background for Circular $\beta$ -Ensembles

### ► Probability density function

$e^{i\theta_1}, \dots, e^{i\theta_n}$  : eigenvalues of Haar-invariant unitary matrix.

pdf:  $f(\theta_1, \dots, \theta_n | \beta = 2)$ , where

$$f(\theta_1, \dots, \theta_n | \beta) = \text{Const} \prod_{1 \leq j < k \leq n} |j e^{i\theta_j} - k e^{i\theta_k}|^\beta$$

$\beta > 0, \theta_i \in [0, 2\pi)$

- This model: *circular  $\beta$ -ensemble* ( $\beta = 1, 2, 4$ ) by physicist Dyson for study of nuclear scattering data



► Three Important Circular Ensembles

COE ( $\beta = 1$ ), CUE ( $\beta = 2$ ), CSE ( $\beta = 4$ )

## ► Three Important Circular Ensembles

COE ( $\beta = 1$ ), CUE ( $\beta = 2$ ), CSE ( $\beta = 4$ )

Construction of COE and CUE

$U = U_{n \times n}$  : Haar unitary

## ► Three Important Circular Ensembles

COE ( $\beta = 1$ ), CUE ( $\beta = 2$ ), CSE ( $\beta = 4$ )

Construction of COE and CUE

$U = U_{n \times n}$  : Haar unitary

- $U$  follows CUE

## ► Three Important Circular Ensembles

COE ( $\beta = 1$ ), CUE ( $\beta = 2$ ), CSE ( $\beta = 4$ )

Construction of COE and CUE

$U = U_{n \times n}$  : Haar unitary

- $U$  follows CUE
- $U^T U$  follows COE

## ► Three Important Circular Ensembles

COE ( $\beta = 1$ ), CUE ( $\beta = 2$ ), CSE ( $\beta = 4$ )

Construction of COE and CUE

$U = U_{n \times n}$  : Haar unitary

- $U$  follows CUE
- $U^T U$  follows COE
- CSE is similar but a bit involved (see Mehta)

## ► Three Important Circular Ensembles

COE ( $\beta = 1$ ), CUE ( $\beta = 2$ ), CSE ( $\beta = 4$ )

Construction of COE and CUE

$U = U_{n \times n}$  : Haar unitary

- $U$  follows CUE
- $U^T U$  follows COE
- CSE is similar but a bit involved (see Mehta)

Entries of  $CUE$  : roughly independent  $\mathbb{C}N(0, 1)$  (Jiang, AP06)

## ► Three Important Circular Ensembles

COE ( $\beta = 1$ ), CUE ( $\beta = 2$ ), CSE ( $\beta = 4$ )

Construction of COE and CUE

$U = U_{n \times n}$  : Haar unitary

- $U$  follows CUE
- $U^T U$  follows COE
- CSE is similar but a bit involved (see Mehta)

Entries of  $CUE$  : roughly independent  $\mathbb{C}N(0, 1)$  (Jiang, AP06)

Entries of  $COE$  : roughly  $\mathbb{C}N(0, 1)$  (but dependent) (Jiang, JMP09)

## ► Three Important Circular Ensembles

COE ( $\beta = 1$ ), CUE ( $\beta = 2$ ), CSE ( $\beta = 4$ )

Construction of COE and CUE

$U = U_{n \times n}$  : Haar unitary

- $U$  follows CUE
- $U^T U$  follows COE
- CSE is similar but a bit involved (see Mehta)

Entries of *CUE* : roughly independent  $\mathbb{C}N(0, 1)$  (Jiang, AP06)

Entries of *COE* : roughly  $\mathbb{C}N(0, 1)$  (but dependent) (Jiang, JMP09)

Killip & Nenciu: Matrix models for circular ensembles



# Moments for Circular $\beta$ -Ensembles

# Moments for Circular $\beta$ -Ensembles

► Bad news from COE:

# Moments for Circular $\beta$ -Ensembles

- ▶ **Bad news from COE:** Let  $M_n$  be COE. By elementary check

$$\mathbb{E}[\text{Tr}(M_n)^2] = \frac{2n}{n+1}$$

# Moments for Circular $\beta$ -Ensembles

- ▶ **Bad news from COE:** Let  $M_n$  be COE. By elementary check

$$\mathbb{E}[\text{Tr}(M_n)^2] = \frac{2n}{n+1}$$

- Moments depend on  $n$

# Moments for Circular $\beta$ -Ensembles

- **Bad news from COE:** Let  $M_n$  be COE. By elementary check

$$\mathbb{E}[\text{Tr}(M_n)^2] = \frac{2n}{n+1}$$

- Moments depend on  $n$
- Later results:  $\mathbb{E}[\text{Tr}(M_n)^2]$  not depend on  $n$  only at  $\beta = 2$

# Moments for Circular $\beta$ -Ensembles

► **Bad news from COE:** Let  $M_n$  be COE. By elementary check

$$\mathbb{E}[\text{Tr}(M_n)^2] = \frac{2n}{n+1}$$

- Moments depend on  $n$
- Later results:  $\mathbb{E}[\text{Tr}(M_n)^2]$  not depend on  $n$  only at  $\beta = 2$
- This suggest: moments for general  $\beta$ -ensemble depend on  $n$

## ► Notation

- $\lambda = (\lambda_1, \lambda_2, \dots)$  : *partition*

## ► Notation

- $\lambda = (\lambda_1, \lambda_2, \dots)$  : *partition*
- $|\lambda| = \lambda_1 + \lambda_2 + \dots$  : *weight*



## ► Notation

- $\lambda = (\lambda_1, \lambda_2, \dots)$  : *partition*
- $|\lambda| = \lambda_1 + \lambda_2 + \dots$  : *weight*
- $m_i(\lambda)$  : *multi of  $i$  in  $(\lambda_1, \lambda_2, \dots)$*

## ► Notation

- $\lambda = (\lambda_1, \lambda_2, \dots)$  : *partition*
- $j\lambda_j = \lambda_1 + \lambda_2 + \dots$  : *weight*
- $m_i(\lambda)$  : *multi of  $i$  in  $(\lambda_1, \lambda_2, \dots)$*
- $l(\lambda) = \#$  of positive  $\lambda_i$  in  $\lambda$  : *length*

## ► Notation

- $\lambda = (\lambda_1, \lambda_2, \dots)$  : *partition*
- $j\lambda_j = \lambda_1 + \lambda_2 + \dots$  : *weight*
- $m_i(\lambda)$  : *multi of  $i$  in  $(\lambda_1, \lambda_2, \dots)$*
- $l(\lambda) = \#$  of positive  $\lambda_i$  in  $\lambda$  : *length*

$$z_\lambda = \prod_{i \geq 1} i^{m_i(\lambda)} m_i(\lambda)!$$

## ► Notation

- $\lambda = (\lambda_1, \lambda_2, \dots)$  : *partition*
- $j\lambda_j = \lambda_1 + \lambda_2 + \dots$  : *weight*
- $m_i(\lambda)$  : *multi of  $i$  in  $(\lambda_1, \lambda_2, \dots)$*
- $l(\lambda) = \#$  of positive  $\lambda_i$  in  $\lambda$  : *length*

$$z_\lambda = \prod_{i \geq 1} i^{m_i(\lambda)} m_i(\lambda)!$$

- $p_\lambda = \prod_{i=1}^{l(\lambda)} p_{\lambda_i}$ , where  $p_k(x_1, x_2, \dots) = x_1^k + x_2^k + \dots$

## ► Notation

- $\lambda = (\lambda_1, \lambda_2, \dots)$  : *partition*
- $j\lambda_j = \lambda_1 + \lambda_2 + \dots$  : *weight*
- $m_i(\lambda)$  : *multi of  $i$  in  $(\lambda_1, \lambda_2, \dots)$*
- $l(\lambda) = \#$  of positive  $\lambda_i$  in  $\lambda$  : *length*

$$z_\lambda = \prod_{i \geq 1} i^{m_i(\lambda)} m_i(\lambda)!$$

- $p_\lambda = \prod_{i=1}^{l(\lambda)} p_{\lambda_i}$ , where  $p_k(x_1, x_2, \dots) = x_1^k + x_2^k + \dots$

$$\lambda = (3, 2, 2) : j\lambda_j = 7, m_2(\lambda) = 2, m_3(\lambda) = 1, l(\lambda) = 3,$$

$$p_\lambda = (\sum_i \lambda_i^3) (\sum_i \lambda_i^2)^2$$

$\alpha > 0, K \geq 1, n \geq 1$ , define

$$A = \left(1 - \frac{j\alpha}{n} \frac{1j}{K + \alpha} \delta(\alpha \geq 1)\right)^K$$

$$B = \left(1 + \frac{j\alpha}{n} \frac{1j}{K + \alpha} \delta(\alpha < 1)\right)^K$$

$\alpha > 0, K \geq 1, n \geq 1$ , define

$$A = \left(1 - \frac{j\alpha}{n} \frac{1j}{K + \alpha} \delta(\alpha \geq 1)\right)^K$$

$$B = \left(1 + \frac{j\alpha}{n} \frac{1j}{K + \alpha} \delta(\alpha < 1)\right)^K$$

Let  $\theta_1, \dots, \theta_n \stackrel{i.i.d.}{\sim} f(\theta), \alpha = 2/\beta$ .

- $Z_n = (e^{i\theta_1}, \dots, e^{i\theta_n})$ ,
- $p_\mu(Z_n) = p_\mu(e^{i\theta_1}, \dots, e^{i\theta_n})$

## Theorem

(a) If  $n = j\mu$ , then

$$A \frac{\mathbb{E}[j p_{\mu}(Z_n) j^2]}{\alpha^{l(\mu)} z_{\mu}} B$$



## Theorem

(a) If  $n \in K = j\mu j$ , then

$$A \quad \frac{\mathbb{E}[j p_\mu(Z_n) j^2]}{\alpha^{l(\mu)} z_\mu} \quad B$$

(b) If  $j\mu j \notin j\nu j$ , then  $\mathbb{E}[p_\mu(Z_n) \overline{p_\nu(Z_n)}] = 0$ .

## Theorem

(a) If  $n \in K = j\mu j$ , then

$$A \frac{\mathbb{E}[j p_\mu(Z_n) j^2]}{\alpha^{l(\mu)} z_\mu} B$$

(b) If  $j\mu j \notin j\nu j$ , then  $\mathbb{E}[p_\mu(Z_n) \overline{p_\nu(Z_n)}] = 0$ .

If  $\mu \notin \nu$  and  $n \in K = j\mu j \cup j\nu j$ , then

$$\left| \mathbb{E}[p_\mu(Z_n) \overline{p_\nu(Z_n)}] \right| \leq \max\{j_A, j_B\} \alpha^{(l(\mu)+l(\nu))/2} (z_\mu z_\nu)^{1/2}$$

## Theorem

(a) If  $n \in K = j\mu j$ , then

$$A \frac{\mathbb{E}[jp_\mu(Z_n)f^2]}{\alpha^{l(\mu)}z_\mu} B$$

(b) If  $j\mu j \notin j\nu j$ , then  $\mathbb{E}[p_\mu(Z_n)\overline{p_\nu(Z_n)}] = 0$ .

If  $\mu \notin \nu$  and  $n \in K = j\mu j \cup j\nu j$ , then

$$\left| \mathbb{E}[p_\mu(Z_n)\overline{p_\nu(Z_n)}] \right| \leq \max\{j_A, j_B\} \alpha^{(l(\mu)+l(\nu))/2} (z_\mu z_\nu)^{1/2}$$

(c)  $\exists C = C(\beta)$  s.t.  $\forall m \geq 1, n \geq 2$

$$\left| \mathbb{E}[jp_m(Z_n)f^2] \right| \leq C \frac{n^3 2^{n\beta}}{m^{1 \wedge \beta}}$$

Take  $\beta = 2$ , then  $A = B = 1$ . We recover

► Theorem (Diaconis and Evans: 2001)

$a = (a_1, \dots, a_k)$ ,  $b = (b_1, \dots, b_k)$  with  $a_j, b_j \geq 0, 1, 2, \dots, g$ .

For  $n = \sum_{j=1}^k j a_j = \sum_{j=1}^k j b_j$ ,

$$\mathbb{E} \left[ \prod_{j=1}^k (\text{Tr}(U_n^j))^{a_j} \overline{(\text{Tr}(U_n^j))^{b_j}} \right] = \delta_{ab} \prod_{j=1}^k j^{a_j} a_j!$$

## Corollary

$\delta\beta > 0$ ,

$$(a) \quad \lim_{n \rightarrow \infty} \mathbb{E} \left[ p_\mu(Z_n) \overline{p_\nu(Z_n)} \right] = \delta_{\mu\nu} \left( \frac{2}{\beta} \right)^{l(\mu)} z_\mu;$$

## Corollary

$\delta\beta > 0$ ,

$$(a) \quad \lim_{n \rightarrow \infty} \mathbb{E} \left[ p_\mu(Z_n) \overline{p_\nu(Z_n)} \right] = \delta_{\mu\nu} \left( \frac{2}{\beta} \right)^{l(\mu)} z_\mu;$$

$$(b) \quad \lim_{m \rightarrow \infty} \mathbb{E} [j p_m(Z_n) f^2] = n \quad \text{for any } n \geq 2.$$

## Corollary

$\mu \neq \nu : K = j\mu j - j\nu j$ . If  $n \geq 2K$ , then

$$(a) \quad \left| \frac{\mathbb{E}[j p_\mu(Z_n) j^2]}{\alpha^{l(\mu)} z_\mu} - 1 \right| \leq \frac{6j1 \alpha j K}{n};$$

## Corollary

$\mu \neq \nu : K = j_{\mu} j_{\nu}$ . If  $n \geq 2K$ , then

$$(a) \quad \left| \frac{\mathbb{E}[j p_{\mu}(Z_n) j^2]}{\alpha^{l(\mu)} z_{\mu}} - 1 \right| \leq \frac{6j^2 \alpha^{jK}}{n};$$

$$(b) \quad \left| \mathbb{E} \left[ p_{\mu}(Z_n) \overline{p_{\nu}(Z_n)} \right] \right| \leq \frac{6j^2 \alpha^{jK}}{n} \alpha^{(l(\mu)+l(\nu))/2} (z_{\mu} z_{\nu})^{1/2}.$$



► Exact formula

The exact formula gives

$$\mathbb{E}[jp_1(Z_n)f^2] = \frac{2}{\beta n} \frac{n}{1 + 2\beta^{-1}}$$

► Exact formula

The exact formula gives

$$\mathbb{E}[jp_1(Z_n)f^2] = \frac{2}{\beta} \frac{n}{n(1 + 2\beta^{-1})} = \begin{cases} \frac{2n}{n+1}, & \text{if } \beta = 1 \\ 1, & \text{if } \beta = 2 \\ \frac{n}{2n-1}, & \text{if } \beta = 4 \end{cases}$$

► Exact formula

The exact formula gives

$$\mathbb{E}[jp_1(Z_n)f^2] = \frac{2}{\beta} \frac{n}{n+1} \frac{1}{1+2\beta^{-1}} = \begin{cases} \frac{2n}{n+1}, & \text{if } \beta = 1 \\ 1, & \text{if } \beta = 2 \\ \frac{n}{2n-1}, & \text{if } \beta = 4 \end{cases}$$

Exact formula is given next

## ▶ Jack Polynomial

Jack polynomial  $J_{\lambda}^{(\alpha)} = J_{\lambda}^{(\alpha)}(x_1, \dots, x_n)$  is symmetric in  $x_1, \dots, x_n$

## ▶ Jack Polynomial

Jack polynomial  $J_{\lambda}^{(\alpha)} = J_{\lambda}^{(\alpha)}(x_1, \dots, x_n)$  is symmetric in  $x_1, \dots, x_n$

- $\alpha = 1$ , it is Schur polynomial

## ▶ Jack Polynomial

Jack polynomial  $J_{\lambda}^{(\alpha)} = J_{\lambda}^{(\alpha)}(x_1, \dots, x_n)$  is symmetric in  $x_1, \dots, x_n$

- $\alpha = 1$ , it is Schur polynomial
- $\alpha = 2$ , it is Zonal polynomial

## ▶ Jack Polynomial

Jack polynomial  $J_{\lambda}^{(\alpha)} = J_{\lambda}^{(\alpha)}(x_1, \dots, x_n)$  is symmetric in  $x_1, \dots, x_n$

- $\alpha = 1$ , it is Schur polynomial
- $\alpha = 2$ , it is Zonal polynomial
- $\alpha = 1/2$ , it is Zonal spherical function

## ▶ Jack Polynomial

Jack polynomial  $J_{\lambda}^{(\alpha)} = J_{\lambda}^{(\alpha)}(x_1, \dots, x_n)$  is symmetric in  $x_1, \dots, x_n$

- $\alpha = 1$ , it is Schur polynomial
- $\alpha = 2$ , it is Zonal polynomial
- $\alpha = 1/2$ , it is Zonal spherical function

Orthogonal property:  $Z_n = (e^{i\theta_1}, \dots, e^{i\theta_n})$



# Proofs by Jack Polynomial

## ► Jack Polynomial

Jack polynomial  $J_{\lambda}^{(\alpha)} = J_{\lambda}^{(\alpha)}(x_1$

Write

$$J_{\lambda}^{(\alpha)} = \sum_{\rho: |\rho|=|\lambda|} \theta_{\rho}^{\lambda}(\alpha) p_{\rho}$$
$$p_{\rho} = \sum_{\lambda: |\lambda|=|\rho|} \Theta_{\rho}^{\lambda}(\alpha) J_{\lambda}^{(\alpha)}$$

Write

$$J_{\lambda}^{(\alpha)} = \sum_{\rho: |\rho|=|\lambda|} \theta_{\rho}^{\lambda}(\alpha) p_{\rho}$$
$$p_{\rho} = \sum_{\lambda: |\lambda|=|\rho|} \Theta_{\rho}^{\lambda}(\alpha) J_{\lambda}^{(\alpha)}$$

For  $j_{\mu} j_{\nu} = j_{\nu} j_{\mu} = K$ ,

$$\mathbb{E} \left[ p_{\mu}(Z_n) \overline{p_{\nu}(Z_n)} \right] = \sum_{\lambda \vdash K: l(\lambda) \leq n} \Theta_{\mu}^{\lambda}(\alpha) \Theta_{\nu}^{\lambda}(\alpha) \mathbb{E} (J_{\lambda}^{(\alpha)} \overline{J_{\lambda}^{(\alpha)}})$$

## Use

- explicit form of  $\mathbb{E}(J_\lambda^{(\alpha)} \overline{J_\lambda^{(\alpha)}})$
- relationship between  $\theta_\rho^\lambda(\alpha)$  and  $\Theta_\rho^\lambda(\alpha)$

Use

- explicit form of  $\mathbb{E}(J_\lambda^{(\alpha)} \overline{J_\lambda^{(\alpha)}})$
- relationship between  $\theta_\rho^\lambda(\alpha)$  and  $\Theta_\rho^\lambda(\alpha)$

we have

$$\begin{aligned} & \mathbb{E} \left[ p_\mu(Z_n) \overline{p_\nu(Z_n)} \right] \\ = & \alpha^{l(\mu)+l(\nu)} z_\mu z_\nu \sum_{\lambda \vdash K: l(\lambda) \leq n} \frac{\theta_\mu^\lambda(\alpha) \theta_\nu^\lambda(\alpha)}{C_\lambda(\alpha)} N_\lambda^\alpha(n) \end{aligned}$$

Use

- explicit form of  $\mathbb{E}(J_\lambda^{(\alpha)} \overline{J_\lambda^{(\alpha)}})$
- relationship between  $\theta_\rho^\lambda(\alpha)$  and  $\Theta_\rho^\lambda(\alpha)$

we have

$$\begin{aligned} & \mathbb{E} \left[ p_\mu(Z_n) \overline{p_\nu(Z_n)} \right] \\ &= \alpha^{l(\mu)+l(\nu)} z_\mu z_\nu \sum_{\lambda \vdash K: l(\lambda) \leq n} \frac{\theta_\mu^\lambda(\alpha) \theta_\nu^\lambda(\alpha)}{C_\lambda(\alpha)} N_\lambda^\alpha(n) \end{aligned}$$

$$C_\lambda(\alpha) = \prod_{(i,j) \in \lambda} \left\{ (\alpha(\lambda_i - j) + \lambda'_j - i + 1)(\alpha(\lambda_i - j) + \lambda'_j - i + \alpha) \right\}$$

Use

- explicit form of  $\mathbb{E}(J_\lambda^{(\alpha)} \overline{J_\lambda^{(\alpha)}})$
- relationship between  $\theta_\rho^\lambda(\alpha)$  and  $\Theta_\rho^\lambda(\alpha)$

we have

$$\begin{aligned} & \mathbb{E} \left[ p_\mu(Z_n) \overline{p_\nu(Z_n)} \right] \\ &= \alpha^{l(\mu)+l(\nu)} z_\mu z_\nu \sum_{\lambda \vdash K: l(\lambda) \leq n} \frac{\theta_\mu^\lambda(\alpha) \theta_\nu^\lambda(\alpha)}{C_\lambda(\alpha)} N_\lambda^\alpha(n) \end{aligned}$$

$$C_\lambda(\alpha) = \prod_{(i,j) \in \lambda} \left\{ (\alpha(\lambda_i - j) + \lambda'_j - i + 1)(\alpha(\lambda_i - j) + \lambda'_j - i + \alpha) \right\}$$

$$N_\lambda^\alpha(n) = \prod_{(i,j) \in \lambda} \frac{n + (j - 1)\alpha}{n + j\alpha} \frac{(i - 1)}{i}$$

*Young diagram*

Main proof:

- play  $C_\lambda(\alpha)$
- play  $N_\lambda^\alpha(n)$
- use orthogonal relations of  $\theta_\mu^\lambda(\alpha)$



► Examples

$$\mathbb{E}[jp_1(Z_n)J^4] = \frac{2n\alpha^2(n^2 + 2(\alpha - 1)n - \alpha)}{(n + \alpha - 1)(n + \alpha - 2)(n + 2\alpha - 1)}$$

► Examples

$$\begin{aligned}\mathbb{E}[jp_1(Z_n)J^4] &= \frac{2n\alpha^2(n^2 + 2(\alpha - 1)n - \alpha)}{(n + \alpha - 1)(n + \alpha - 2)(n + 2\alpha - 1)} \\ &= \begin{cases} \frac{8(n^2+2n-2)}{(n+1)(n+3)}, & \text{if } \beta = 1 \\ 2, & \text{if } \beta = 2 \\ \frac{2n^2-2n-1}{(2n-1)(2n-3)}, & \text{if } \beta = 4 \end{cases}\end{aligned}$$

$$\mathbb{E} \left[ p_2(Z_n) \overline{p_1(Z_n)^2} \right]$$

$$\begin{aligned}
& \mathbb{E} \left[ p_2(Z_n) \overline{p_1(Z_n)^2} \right] \\
= & \frac{2\alpha^2(\alpha - 1)n}{(n + \alpha - 1)(n + 2\alpha - 1)(n + \alpha - 2)}
\end{aligned}$$

$$\begin{aligned}
& \mathbb{E} \left[ p_2(Z_n) \overline{p_1(Z_n)^2} \right] \\
&= \frac{2\alpha^2(\alpha - 1)n}{(n + \alpha - 1)(n + 2\alpha - 1)(n + \alpha - 2)} \\
&= \begin{cases} \frac{8}{(n+1)(n+3)}, & \text{if } \beta = 1 \\ 0, & \text{if } \beta = 2 \\ \frac{-1}{(2n-1)(2n-3)}, & \text{if } \beta = 4 \end{cases}
\end{aligned}$$

**The End!**

**Thanks for your patience!**