



# Modeling age-related growth data by a functional Gaussian process

Peter Hall,

*University of California, Davis, USA, and University of Melbourne, Australia*

Hans-Georg Müller

*University of California, Davis, USA*

and Fang Yao

*University of Toronto, Canada*

[Received April 2006. Final revision December 2007]

**S**ummary. In longitudinal data analysis one frequently encounters non-Gaussian data that are repeatedly collected for a sample of individuals over time. The repeated observations could be binomial, Poisson or of another discrete type or could be continuous. The timings of the repeated measurements are often sparse and irregular. We introduce a latent Gaussian process model for such data, establishing a connection to functional data analysis. The functional methods proposed are non-parametric and computationally straightforward as they do not involve a likelihood. We develop functional principal components analysis for this situation and demonstrate the prediction of individual trajectories from sparse observations. This method can handle missing data and leads to predictions of the functional principal component scores which serve as random effects in this model. These scores can then be used for further statistical analysis, such as inference, regression, discriminant analysis or clustering. We illustrate these non-parametric methods with longitudinal data on primary biliary cirrhosis and show in simulations that they are competitive in comparisons with generalized estimating equations and generalized linear mixed models.

**Keywords:** Binomial data; Eigenfunction; Functional data analysis; Functional principal component; Prediction; Random effect; Repeated measurements; Smoothing; Stochastic process

## 1. Introduction

### 1.1. Preliminaries

When undertaking prediction in longitudinal data analysis, in order to give a well-paced and informed measurement, a reliable and informative set of observations is required for each subject, over time and irregular measurements. Irregularity of measurements for individual subjects is an inherent difficulty of such data. The effective use of such data is especially important when the information can be accessed. This is the case for the model here, which has been measured on a set of individuals. We aim at a flexible non-parametric functional data analysis approach, which is in contrast to the commonly used parametric model, such as generalized linear mixed model (GLMM) or generalized estimating equations.

*Address for correspondence:* Hans-Georg Müller, Department of Statistics, University of California at Davis, One Shields Avenue, Davis, CA 95616, USA.  
E-mail: mueller@wald.ucdavis.edu

(GEE) (see, for example, Heagerty (1999) for recent discussion on applying such models to repeated binary measurements, Po Ahmadi (2000) for a related aspect of covariance modelling and Heagerty and Zegeer (2000), Heagerty and Kurland (2001) and Chio and Mullah (2005) for discussion on limitations, modification and feasibility of handling parametric assumptions).

A non-parametric functional approach for the analysis of longitudinal data, in which philosophy of the data is a peak for hemoglobin and in the end flexible, is a special type of model better than the parametric GEE or GLMM approaches in many situations. However, it faces difficulties due to the potential large gap between repeated measurements in typical parametric longitudinal data. The parametric method often comes with a large bias in the parametric form of the handling function. In contrast, in the presence of such gaps, the classical non-parametric approach of smoothing individual observations in a flexible, non-feasible (Yao *et al.*, 2005). The problem has a reduced bias and is accepted in the common endogenous case of non-Gaussian longitudinal responses such as binomial or Poisson responses (see Section 5).

We demonstrate how one can overcome the difficulties that are posed by such data for non-parametric approaches, by applying a modified method of functional data analysis. Functional data analysis methods have been primarily developed for smoothing and denoising sampled data (Ramakrishnan and Siliveru, 2002, 2005). The basic idea of connecting data has a hierarchical analysis of functional data analysis methodology in population longitudinal data in Gaussian processes (LGP) (for example, of latent processes modelling longitudinal disease counts, for example, Diggle *et al.* (1998), Johndrow and Sathar (2002), Hahemi *et al.* (2003) and Poon *et al.* (2006)). Specifically, the Gaussian process makes it possible to overcome the bias condition of a generalised likelihood approach of the hierarchical longitudinal data analysis. The effect of the mean and covariance process of the LGP. Simulation indicates that the method is in practice in the longitudinal data analysis of the Gaussian process.

Since efficient flexible parametric estimation of the longitudinal Gaussian process is often difficult from a large number of parameters, making computation of the maximum likelihood approaches computationally demanding and unstable, especially in the endogenous LGP or random observations for individual observations, a direct method of the link function. The effective specific observations are pooled to the probability of a response in the binary response case. When the link function is a known mean and covariance of the Gaussian process, a method of the unknown maximum likelihood estimation is a good choice of flexibility, but it is a challenging problem of constructing appropriate estimators.

The methodology proposed is a flexible empirical end functional data analysis technology of the case of non-Gaussian repeated measurements. Prominent examples of such data are repeated binary measurements on repeated counts. The method proposed is a model-based Bayesian approach: the estimation of random coefficients may be eliminated, and in this case a simple Taylor approximation may be applied, explicit and non-parametric mean and covariance functions; and the elements of the component are specified of the hierarchical estimation method. The implementation of the estimation has a proposed approach of the flexibility and numerical simplicity.

The analysis of continuous Gaussian parametric longitudinal data by functional methods has been considered previously (e.g. Shi *et al.* (1996), Rice and Wu (2000), James *et al.* (2001) and James and Sgambati (2003)). Our main goal from functional data analysis is functional principal component analysis, the observations are decomposed into a mean function and eigenfunctions (e.g. Rice and Siliveru (1991) and Boente and Faiman (2000)). Various aspects of the hierarchical approach have been functional and longitudinal data analysis discussed in Sani, Ali and Lee (1998), Rice (2004) and Zhao *et al.* (2004); an excellent survey of modelling longitudinal

ajec o ie in biological applica ion i h FPC i Ki kpa ick and Heckman (1989). FPC anal i allo o achie e h ee majo goal:

- (a) dimen ion ed c ion of f nc ional da a b mma i ing he da a in a fe FPC ;
- (b) he p edic ion of indi id al ajec o ie f om pa e da a, b e ima ing he FPC co e of he ajec o ie ;
- (c) f he a i ical anal i of longi dinal da a ba ed on he FPC co e .

In he ne b ec ion, e in od ce he LGP model; hen in Sec ion 2 he p o o ed e i ma e, follo ed b applica ion o p edic ion (Sec ion 3). The e l f om a im la ion d , incl ding a compa ion of he me hod p o o ed i h GLMM and GEE, a e p o ed in Sec ion 4. The anal i of non-Ga ian pa e longi dinal da a i ill a ed in Sec ion 5, i h he longi dinal anal i of he occ rance of hepa omegal in p ima biliar ci ho i. Thi i follo ed b a bief di c ion (Sec ion 6) and an appen i, hich con ain de i a ion and ome heo e ical e l abo e ima ion.

1.2. Latent Gaussian process model

Gene all, deno ing he gene ali ed e pon e b  $Y_{ij}$ , e ob e e independent copie of  $Y$ , b , in each ca e, onl fo a fe pa e ime poin . In pa ic la, he da a e pai  $(T_{ij}, Y_{ij})$ , fo  $1 \leq i \leq n$  and  $1 \leq j \leq m_i$ , he e  $Y_{ij} = Y_i(T_{ij})$  fo an nde l ing r andom ajec o  $Y_i$ , and each  $T_{ij} \in \mathcal{I} = [0, 1]$ . The pa e and ca e ed na e of he ob e a ion ime  $T_{ij}$  ma be e p e ed heo e ical b no ing ha he  $m_i$  a e nifo ml bon ded, if he e an i e ha e a de e min i ic o i gin, o ha he e p e en he al e of independent and iden ical di ib ed andom a iable i h fficien l igh ail, if he  $m_i$  o i gina e o cha ical . We a e aiming a he eem ingl diffic l a k of making ch pa e de ign amenable o f nc ional me hod, hich ha e been p ima il aimed a den el collec ed moo h da a.

A cen al a mp ion fo o app oach i ha he dependence be een he ob e a ion  $Y_{ij}$  i inhe id f om an nde l ing nob e ed Ga ian p o ce  $X$ : le  $Y(t)$ , fo  $t \in \mathcal{T}$ , he e  $\mathcal{T}$  a compac in e al, deno e a o cha ic p o ce a i f ing

$$E\{Y(t_1) \dots Y(t_m) | X\} = \prod_{j=1}^m g\{X(t_j)\}, \tag{1}$$

$$E\{Y(t)^2 | X\} \leq g_1\{X(t)\}$$

fo  $0 \leq t_1 < \dots < t_m \leq 1$  and  $0 < t < 1$ . He e,  $X$  deno e a Ga ian p o ce on  $\mathcal{I}$ ,  $g$  i a moo h, mono one inc ea ing link f nc ion, f om he eal line o he r ange of he di ib ion of he  $Y_{ij}$ , and  $g_1$  i a bon ded f nc ion. Al ho gh e ob e e independent copie of  $Y$ , he e a e acce ible onl fo a fe pa e ime poin fo each bjec . The Ga ian p o ce e  $X_i$  and mea emen ime  $T_{ij}$ , fo  $1 \leq i \leq n$  and  $1 \leq j \leq m_i$ , a e a med o be o all independent, he  $T_{ij}$  a e aken o be iden ical di ib ed a  $\mathcal{T}$ , a , i h p o  $\mathcal{I}$  and he  $X_i$  a e p o ed o be iden ical di ib ed a  $X$ . When in e p e ed fo he da a  $(T_{ij}, Y_{ij})$ , model (1) implie ha

$$E\{Y_i(T_{i1}) \dots Y_i(T_{im_i}) | X_i(T_{i1}), \dots, X_i(T_{im_i})\} = \prod_{j=1}^{m_i} g\{X_i(T_{ij})\}. \tag{2}$$

The a mp ion ha  $X$  a model (1) i Ga ian p o ide a pla ible a of linking o cha ic p o e ie of  $Y(t)$  fo al e  $t$  in diffe en pa e of  $\mathcal{I}$ , o ha da a ha a e ob e ed a each ime poin can be e ed fo inf e nce abo f e eal e of  $Y(t)$  fo an p e cific al e of  $t$ . The idea of pooling da a ac o bjec o o e come he pa ene p oblem i mo i a ed a in Yao

et al. (2005). The link function  $g$  is assumed known; for example we might elect the logit link in the binomial case,  $g(x) = \text{exp}(x) / \{1 + \text{exp}(x)\}$ , and the log-link for count data; under some circumstances, the link can also be estimated non-parametrically. An important special case of model (1) is that of binomial processes, i.e.  $0 \leq l_i \leq 1$  data, here the fitted intensities in model (1) simplify to

$$P\{Y(t_1) = l_1, \dots, Y(t_m) = l_m | X\} = \prod_{j=1}^m g\{X(t_j)\}^{l_j} [1 - g\{X(t_j)\}]^{1-l_j}, \tag{3}$$

for all sequences  $l_1, \dots, l_m$  of 0's and 1's. In this case, the link function  $g$  could be chosen as a distribution function and the meteorological population could be an extension of functional data analysis longitudinal binomial data.

**2. Estimating the accuracy of Gaussian approximation**

To fit model (1) or make predictions inferences about the realizations of  $Y(t)$ , we need to estimate the defining characteristics of the process  $X$ , i.e. its mean and covariance structure. In a sense, the distribution of  $Y$  can be completely specified, e.g. in the binomial model (3), one possible approach would be maximum likelihood. This is, however, a difficult proposition in the irregular case, here it would necessitate the specification of a large number of parameters for the intensity mean and covariance, hence in total, a difficult task which can only be overcome by making efficient assumptions, limiting the flexibility of the approach. Moreover, we are considering a non-stationary case, and the number of parameters would need to increase with  $n$ , the sample size. Finally, another major motivation is the end of functional approach to non-Gaussian longitudinal data. To obtain the non-parametric flavor, we prefer not to make strong assumptions than model (1), and in particular do not wish to make the efficiency assumptions that would be necessary to employ maximum likelihood methods.

Our approach is based on the proposition that the realization of  $X_i$  about its mean is relatively small. In particular, we assume that

$$X_i(t) = \mu(t) + \delta Z_i(t), \quad \mu = E(X_i), \tag{4}$$

where  $Z_i$  is a Gaussian process with zero mean and bounded covariance and  $\delta > 0$  is an unknown small constant. In this case, assuming that  $g$  has a bounded derivative, and writing  $(X, Z)$  for a generic pair  $(X_i, Z_i)$ , we have

$$g(X) = g(\mu) + \delta Z g^{(1)}(\mu) + \frac{1}{2} \delta^2 Z^2 g^{(2)}(\mu) + \frac{1}{6} \delta^3 Z^3 g^{(3)}(\mu) + O_p(\delta^4), \tag{5}$$

$$E[g\{X(t)\}] = g(\mu) + \frac{1}{2} \delta^2 E\{Z^2(t)\} g^{(2)}\{\mu(t)\} + O(\delta^4) \tag{6}$$

and

$$\text{cov}[g\{X(s)\}, g\{X(t)\}] = \delta^2 g^{(1)}\{\mu(s)\} g^{(1)}\{\mu(t)\} \text{cov}\{Z(s), Z(t)\} + O(\delta^4). \tag{7}$$

Here and hereafter we make the assumption that  $g^{(1)}$  does not vanish, and has  $\inf_{s \in D} \{g^{(1)}(s)\} > 0$ , hence  $D$  is the (compact) range of the mean function  $\mu$ . Setting

$$\left. \begin{aligned} \alpha(t) &= E[g\{X(t)\}], \\ \nu(t) &= g^{-1}\{\alpha(t)\}, \\ \tau(s, t) &= \text{cov}[g\{X(s)\}, g\{X(t)\}] / g^{(1)}\{\mu(s)\} g^{(1)}\{\mu(t)\}, \end{aligned} \right\} \tag{8}$$

we obtain

$$\mu(t) = E\{X(t)\} = g^{-1}(E[g\{X(t)\}]) + O(\delta^2) = \nu(t) + O(\delta^2), \tag{9}$$

$$\sigma(s, t) = \text{co}\{X(s), X(t)\} = \frac{\text{co}[g\{X(s)\}, g\{X(t)\}]}{g^{(1)}\{\mu(s)\}g^{(1)}\{\mu(t)\}} + O(\delta^4) = \tau(s, t) + O(\delta^4). \tag{10}$$

The e f o r m a l e i m m e d i a t e l y e g g e s e i m a o f  $\mu$  and  $\sigma$ , if e a e i l l i n g o n e g l e c t h e e f f e c t o f o r d e r  $O(\delta^2)$ . I n d e e d, e m a e i m a e

$$\alpha(t) = E\{Y(t)\} = E[E\{Y(t)|X(t)\}] = E[g\{X(t)\}], \tag{11}$$

b p a r a m e t e r s a r e h o l d e r s o f  $(T_{ij}, Y_{ij})$ , and e i m a e

$$\beta(s, t) = E\{Y(s)Y(t)\} = E[g\{X(s)\}g\{X(t)\}] \tag{12}$$

(b e i n g m o d e l (1)) b p a r a m e t e r s a r e h o l d e r s o f  $((T_{ij}, T_{ik}), Y_{ij}Y_{ik})$  f o r  $1 \leq i \leq n$ , w h e r e  $m_i \geq 2$ , and  $1 \leq j, k \leq m_i$  i f  $j \neq k$ . I n n e c e s s a r y c a s e s t h e d i a g o n a l e n t r i e s i n  $h_{ij}$  m o d e l (1) e h a v e

$$E\{Y^2(t)\} = E[E\{Y^2(t)|X(t)\}] > E[E\{Y(t)|X(t)\}]^2 = E[g\{X(t)\}]^2,$$

h e n e e a r e  $\{Y(t)|X(t)\} > 0$ , s o t h e a r i a n c e a l o n g t h e d i a g o n a l i n g e n e r a l i l l h a s a n e i g e n c o m p o n e n t, l e a d i n g t o a c o a r i a n c e s u r f a c e h a s a d i s c o n t i n u i t y a l o n g t h e d i a g o n a l. M o r e d e t a i l s a b o u t t h i s p h e n o m e n o n c a n b e f o u n d i n Y a o *et al.* (2005). I m p l e m e n t a t i o n o f t h e e i m a o f  $h_{ij}$  m o d e l (1) b e i n g l o c a l l e a r n e d e i m a o f  $h_{ij}$  i s d i s c u s s e d i n A p p e n d i x A.

F r o m t h e e i m a o f  $h_{ij}$ ,  $\alpha$  and  $\beta$  o f  $\alpha$  and  $\beta$  e s p e c i e l y, e o b a i n e i m a o f

$$\nu(t) = g^{-1}\{\alpha(t)\}, \tag{13}$$

$$\tau(s, t) = \{\beta(s, t) - \alpha(s)\alpha(t)\} / g^{(1)}\{\nu(s)\}g^{(1)}\{\nu(t)\}$$

f o r

$$\nu(t) = g^{-1}\{\alpha(t)\}, \tag{14}$$

$$\tau(s, t) = \{\beta(s, t) - \alpha(s)\alpha(t)\} / g^{(1)}\{\nu(s)\}g^{(1)}\{\nu(t)\}$$

e s p e c i e l y. B e i n g o f a p p o i m a t i o n (9) and (10) e m a i n e p e  $\nu$  and  $\tau$  a e i m a o f  $\mu$  and  $\sigma$  e s p e c i e l y, i. e. e

$$\mu(t) = \nu(t), \tag{15}$$

$$\sigma(s, t) = \tau(s, t).$$

T h e e i m a o f  $h_{ij}$  d o n o t d e p e n d o n t h e c o n s t a n t  $\delta$ , w h i c h h e r e f o r e d o e s n o t n e e d t o b e k n o w n o r e i m a d. A l t h o u g h t h e e i m a o f  $\tau(s, t)$  i s i m m e d i a t e l y g e n e r a l l y n o e n j o y t h e p o s i t i v e s e m i d e f i n e n e s s p o p e r t y h a s i t e i s e d o f a c o a r i a n c e f u n c t i o n. T h i s d e f i c i e n c y c a n b e o v e r c o m e b y i m p l e m e n t i n g a m e t h o d h a s a d e s c r i b e d i n Y a o *et al.* (2003), w h i c h i s o d e r f o m t h e s p e c i a l d e c o m p o s i t i o n o f  $\tau$  h o e e m a h a c o e p o n d o n e g a t i v e e i g e n a l e s. I n e a c h o f t h e s e c a s e s, i n d o i n g s o, t h e m e a n a e d e o f  $\tau$  i s i m m e d i a t e l y i m p o e d b o m i n g a e m h a c o e p o n d o n a n e g a t i v e e i g e n a l e; d e t a i l s c a n b e f o u n d i n A p p e n d i x B. I n t h e f o l l o w i n g e o r k i n g t h e e i m a o f  $h_{ij}$  a e d e f i n e d i n A p p e n d i x B. P o p e r t y o f t h e e i m a o f  $\alpha$  and  $\beta$ , and  $\nu$  and  $\tau$ , w h i c h a r e d e f i n e d a e p e e i o n (32), (33) and (13) e s p e c i e l y, and o f e i m a o f  $\mu$  and  $\sigma$  a e p e e i o n (15) a e d i s c u s s e d i n A p p e n d i x C.

**3. Predicting individual trajectories of the effect**

**3.1. Predicting functional principal component scores**

One of the main purposes of the functional data analysis model proposed in dimension reduction is to predict FPC scores. The prediction of the underlying hidden Gaussian process for the subject in a study. Specifically, the predicted FPC scores provide a mean for evaluating the individual data, and also for dimension reduction, and can be used for inference, discriminant analysis or regression.

The starting point is the Karhunen-Loève expansion of random process  $X_i$  of the LGP,

$$X_i(t) = \mu(t) + \sum_{j=1}^{\infty} \xi_{ij} \psi_j(t), \tag{16}$$

where  $\psi_j$  are the orthonormal eigenfunctions of the linear integral operator  $B$  with kernel  $\sigma(s, t)$ , having an  $L^2$ -function  $f$  on  $Bf(s) = \int \sigma(s, t) f(t) dt$ , i.e. the eigenfunctions of

$$\int \text{cov}\{X(s), X(t)\} \psi_j(t) ds = \theta_j \psi_j(t),$$

where  $\theta_j$  is the eigenvalue having a associated eigenfunction  $\psi_j$ . The  $\xi_{ij} = \int \{X_i(t) - \mu(t)\} \psi_j(t) dt$  are the FPC scores having the role of random effects, with  $E(\xi_{ij}) = 0$  and  $\text{var}(\xi_{ij}) = \theta_j$ , where  $\theta_j$  is the eigenvalue corresponding to eigenfunction  $\psi_j$ . Once the image of  $\sigma(s, t)$  (15) has been determined, the corresponding image of  $\theta_j$  and  $\psi_j$  of eigenvalue and eigenfunction of the eigenprocess  $X$  are obtained by a standard discrete eigenvalue procedure, where the image are determined from a discrete principal component analysis.

We aim to estimate the linear process

$$E\{X_i(t) | Y_{i1}, \dots, Y_{im}\} = \sum_{j=1}^{\infty} E(\xi_{ij} | Y_{i1}, \dots, Y_{im}) \psi_j(t) \tag{17}$$

of the process  $X_i$ , given the data  $Y_{i1}, \dots, Y_{im}$ . Here a reduction of the expansion is needed only the first  $M$  components are needed. Then, focusing on the first  $M$  conditional FPC scores will allow us to reduce the dimension of the problem and also to evaluate the high frequency data. According to equation (17), the task of estimation and prediction of individual processes can be reduced to that of estimating  $E(\xi_{ij} | Y_{i1}, \dots, Y_{im})$ . In the following we develop a simple approximation in the non-Gaussian case by means of a moment-based approach, as follows. The repeated measurements process is assumed to be generated by

$$Y_{ik} = Y_i(T_{ik}) = g\{X_i(T_{ik})\} + e_{ik}, \tag{18}$$

with independent errors  $e_{ik}$ , assuming

$$\begin{aligned} E(e_{ik}) &= 0, \\ \text{var}(e_{ik}) &= \gamma^2 v\{g\{X_i(T_{ik})\}\}. \end{aligned} \tag{19}$$

Here,  $\gamma^2$  is an unknown variance (or dispersion) parameter and  $v(\cdot)$  is a known smooth variance function, which is determined by the characteristics of the data. For example, in the case of repeated binary observations, one could choose  $v(u) = u(1-u)$ . In the following we implicitly condition on the measurement times  $T_{ij}$ .

With a Taylor expansion of  $g$ , using the expansion (4) and assuming a before hand  $\inf\{g^{(1)}(\cdot)\} > 0$ , we obtain

$$g\{X(t)\} = g\{\mu(t)\} + g^{(1)}\{\mu(t)\}\{X(t) - \mu(t)\} + O(\delta^2). \tag{20}$$

Defining

$$\varepsilon_{ik} = \frac{e_{ik}}{g^{(1)}\{\mu(T_{ik})\}},$$

$$U_{ik} = \mu(T_{ik}) + \frac{Y_{ik} - g\{\mu(T_{ik})\}}{g^{(1)}\{\mu(T_{ik})\}},$$

equations (19) and (20) lead to  $U_{ik} = X_i(T_{ik}) + \varepsilon_{ik} + O(\delta^2)$ . We note also the identity (15) and equation (19) lead to

$$\tilde{e}_{ik} = Z_{ik}\gamma \frac{v[g\{\mu(T_{ik})\}]^{1/2}}{g^{(1)}\{\mu(T_{ik})\}},$$

where the  $Z_{ik}$  are independent copies of a standard Gaussian  $N(0, 1)$  random variable, so that the first-order moments of  $\tilde{e}_{ik}$  are approximating those of  $\varepsilon_{ik}$ . Then, for small  $\delta$ ,  $U_{ik} \approx X_i(T_{ik}) + \tilde{e}_{ik}$ , implying that

$$E(\xi_{ij}|Y_{i1}, \dots, Y_{im_i}) = E(\xi_{ij}|U_{i1}, \dots, U_{im_i}) \approx E\{\xi_{ij}|X_i(T_{i1}) + \tilde{e}_{i1}, \dots, X_i(T_{im_i}) + \tilde{e}_{im_i}\}.$$

Observe that the Gaussian approximation for  $U_{ik}$  in place of  $X_i$ , the latter conditional expectation is seen to be a linear function of the elements on the right-hand side, and hence

$$E(\xi_{ij}|Y_{i1}, \dots, Y_{im_i}) = A_{ij}\tilde{X}_i \tag{21}$$

is a reasonable prediction for the random effect  $\xi_{ij}$ , where  $\tilde{X}_i = (X_i(T_{i1}) + \tilde{e}_{i1}, \dots, X_i(T_{im_i}) + \tilde{e}_{im_i})^T$  and the  $A_{ij}$  are matrices depending only on  $\gamma, \mu, v, g$  and  $g^{(1)}$ . These quantities are either known or estimable, with the sole exception of  $\gamma$ , the estimation of which is discussed below. The explicit form of equation (21) is given in Appendix D.

### 3.2. Predicting trajectories

Moreover, based on equations (16) and (21), prediction is achieved for the LGP, as obtained as

$$X_i(t) = E\{X_i(t)|Y_{i1}, \dots, Y_{im_i}\} = \mu(t) + \sum_{j=1}^M A_{ij}\tilde{X}_i\psi_j(t), \tag{22}$$

and prediction is achieved for the observed process  $Y$  as

$$Y_i(t) = E\{Y_i(t)|Y_{i1}, \dots, Y_{im_i}\} = g\{X_i(t)\}, \tag{23}$$

where  $t$  may be any time point within the range of process  $Y$ , including times for which no response are observed. Predicted values for  $Y(t)$  can, of course, be used to predict the entire response distribution when the mean determines the entire distribution, such as in binomial and Poisson cases. This method could also be employed for the prediction of missing values in a situation where missing data occur on all a random.

To evaluate the effect of a variable on the prediction, we use a cross-sectional comparison where we compare prediction of  $Y_{ik}$ , which are obtained by leaving a observation out, with  $Y_{ik}$  itself. Comparing

$$Y_{ik}^{(-ik)} = E(Y_{ik}|Y_{i1}, \dots, Y_{i,k-1}, Y_{i,k+1}, \dots, Y_{im_i}) = g\{X_i^{(-ik)}(T_{ik})\}, \quad 1 \leq i \leq n, \quad 1 \leq k \leq m_i, \tag{24}$$

where

$$X_i^{(-ik)}(T_{ik}) = \mu(t) + \sum_{j=1}^M E(\xi_{ij} | Y_{i1}, \dots, Y_{i,k-1}, Y_{i,k+1}, \dots, Y_{im_i}) \psi_j(t), \tag{25}$$

we define the Pearson coefficient of determination

$$PE(\gamma^2) = \sum_{i,k} \frac{(Y_{ik}^{(-ik)} - Y_{ik})^2}{v[g\{X_i^{(-ik)}(T_{ik})\}]}, \tag{26}$$

which will depend on the variance parameter  $\gamma^2$  and implicitly also on the number of eigenfunctions  $M$  that are included in the model; see equation (19).

We found that the following iterative selection procedure, for choosing the number of eigenfunctions  $M$  and the optimal variance parameter  $\gamma^2$  in a one-dimensional, led to good practical results: choose a starting value for  $M$ ; then obtain  $\gamma^2$  by minimizing the cross-validated prediction error  $PE$  in the expected  $\gamma^2$ ,

$$\gamma = \text{arg min}_{\gamma} \{PE(\gamma^2)\}. \tag{27}$$

Then, in a bootstrap procedure, predict  $M$  by the criterion has just described below, and repeat the entire process until the values of  $M$  and  $\gamma^2$  stabilize. This iterative algorithm is coded in the following practical algorithm for  $M$  should be 2 or 3.

Specifically, for the choice of  $M$ , we adopt a quasi-likelihood-based functional information criterion FIC that is an extension of the Akaike information criterion AIC for functional data (see Yao *et al.* (2005) for a related pseudo-Gaussian likelihood-based criterion). The number of eigenfunctions  $M$ , to be included in the model, is chosen in such a way as to minimize

$$FIC(M) = -2 \sum_{i,k} \int_{Y_{ik}}^{Y_{ik}} \frac{Y_{ij} - t}{\gamma^2 v(t)} dt + 2M. \tag{28}$$

The penalty  $2M$  corresponding to has just been defined in AIC; otherwise, the penalty corresponding to the Bayesian information criterion BIC could be used as well.

Some implementation issues can be imposed in this criterion for the choice of  $M$  and  $\gamma^2$ , so that loops cannot happen, although the need to be observed here occurs. We also investigated direct minimization of equation (26) in a one-dimensional form for both  $\gamma$  and  $M$ . Besides being computationally more complex in general, this alternative minimization scheme ended up choosing more components and ended in less satisfactory results, especially when observing the prediction error. In each of making a practical comparison about the variance functional  $v$ , in some cases it may be preferable to estimate it non-parametrically. This can be done via empirical quasi-likelihood regression (Chio and Müller, 2005).

## 4. Simulation

### 4.1. Comparisons with generalized estimating equations and generalized linear mixed models

The simulation is based on latent processes  $X(t)$  with mean function  $E\{X(t)\} = \mu(t) = 2 \sin(\pi t/5)/\sqrt{5}$ , and covariance  $\{X(s), X(t)\} = \lambda_1 \phi_1(s) \phi_1(t)$  derived from a single eigenfunction  $\phi_1(t) = -\cos(\pi t/10)/\sqrt{5}$ ,  $0 \leq t \leq 10$ , with eigenvalue  $\lambda_1 = 2$  ( $\lambda_k = 0, k \geq 2$ ). Then 200 Gaussian and 200 non-Gaussian samples of latent processes consisting of  $n = 100$  random observations each are generated by  $X_i(t) = \mu(t) + \xi_{i1} \phi_1(t)$ , where for the 200 Gaussian samples the FPC coefficients  $\xi_{i1}$  are simulated from  $\mathcal{N}(0, 2)$ , whereas for the non-Gaussian samples they are simulated from a mixture of normal distributions:  $\mathcal{N}(\sqrt{2}, 2)$  with probability  $\frac{1}{2}$  and  $\mathcal{N}(-\sqrt{2}, 2)$



ih probabilities  $\frac{1}{2}$ . Binao come  $Y_{ij}$  e, e genera ed a Be no lli a iable ih probabilities  $E\{Y_{ij}|X_i(t_{ij})\} = g\{X_i(t_{ij})\}$ , sing he canonical logi link f nc ion  $g^{-1}(p) = \log\{p/(1-p)\}$  fo  $0 < p < 1$ .

To genera e he pa e ob e a ion, each a jec o a a ppled a a a ndom n mbe of poin, cho en nifo ml f om  $\{8, \dots, 12\}$ , and he loca ion of he mea remen e e nifo ml di ib ed o e he domain  $[0, 10]$ . Fo he moo hing ep, ni a ia e and bi a ia e p od c Epanechniko eigh f nc ion e e ed, i.e.  $K_1(x) = (3/4)(1-x^2) \mathbf{1}_{[-1,1]}(x)$  and  $K_2(x, y) = (9/16)(1-x^2)(1-y^2) \mathbf{1}_{[-1,1]}(x) \mathbf{1}_{[-1,1]}(y)$ , he e  $\mathbf{1}_A(x)$  e al 1 if  $x \in A$  and 0 o he i e fo an e  $A$ . The n mbe of eigen f nc ion  $M$  and he o e di pe ion pa ame e  $\gamma^2$  e e epa a el e lec ed fo each n b he i e a ion (27) and e a ion (28). The e i e a ion con e ged fa e i ing onl 2 4 i e a ion ep in mo ca e.

We compa e he non-pa ame ic LGP me hod p o po ed ih he pop la pa ame ic appoache p o ided b GLMM and GEE. Fo he GEE me hod, e ed he n c ed co elation op ion and boh GEE and GLMM e e n ih linea (me hod GEE-L and GLMM-L) and in addi ion ih ad a ic (me hod GEE-Q and GLMM-Q) fi ed effec. We e fo c i e ia fo he compa ion, mea ing di c epancie be en e ima e and a ge boh in e m of la en p o ce e  $X$  and e pon e p o ce e  $Y = g(X)$ , and compa ing boh e ima e fo mean f nc ion  $\mu = E(X)$  and  $g(\mu)$  e pec i el and p edic ion of b jec p e cific a jec o i e  $X_i$  and  $g(X_i)$  e pec i el. The la e a e a ilable fo he LGP and GLMM me hod, b no fo GEE, hich aim a ma ginal modelling. The p e cific c i e ia fo he compa ion a e a follo :

$$XMSE = \int_{\mathcal{I}} \{\mu(t) - \mu(t)\}^2 dt / \int_{\mathcal{I}} \mu^2(t) dt, \tag{29}$$

$$YMSE = \int_{\mathcal{I}} [g\{\mu(t)\} - g\{\mu(t)\}]^2 dt / \int_{\mathcal{I}} g^2\{\mu(t)\} dt,$$

$$XPE_i = \int_{\mathcal{I}} \{X_i(t) - X_i(t)\}^2 dt / \int_{\mathcal{I}} X_i^2(t) dt, \tag{30}$$

$$YPE_i = \int_{\mathcal{I}} [g\{X_i(t)\} - g\{X_i(t)\}]^2 dt / \int_{\mathcal{I}} g^2\{X_i(t)\} dt,$$

fo  $i = 1, \dots, n$ . S mma a i ic fo he al e of he e c i e ia f om 200 Mon e Ca lo n a e ho n in Table 1.

The e e l indica e ha, fi of all, he LGP me hod p o po ed i no en i i e o he Ga ian a mp ion fo la en p o ce e. Al ho gh he e i ome de e io a ion in he non-Ga ian ca e, i i minimal. Thi non- en i i i o he Ga ian a mp ion ha been de c ibed befo e in f nc iona da a nal i in he con e of p nc ipal anal i b condi onal e pec a ion (ee Yao *et al.* (2005)). Secondl, he non-linea i in he a ge f nc ion ho he pa ame ic me hod off ack, e en hen he mo e fle ible ad a ic fi ed effec e ion a e ed. We find ha he LGP me hod con e clea ad an age in e ima ion and e peciall in p edic ing indi id a jec o i e in ch i a ion. Whe ea he pa ame ic me hod a e en i i e o iola ion of a mp ion, he LGP me hod i de igned o o k nde minimal a mp ion and he e fo e p o ide a ef l al e na i e appoach.

4.2. Effect of the size of variation

He e e e amine he infl ence of he i e of he a ia ion con an  $\delta$  on model e ima ion, incl ding mean f nc ion, eigen f nc ion and indi id al a jec o i e. In addi ion o c i e ia (29)

**Tab e 1.** Simulation results for the comparisons of mean estimates and individual trajectory predictions obtained by the proposed non-parametric LGP method with those obtained for the established parametric methods GLMM-L, GLMM-Q, GEE-L and GEE-Q, with linear and quadratic fixed effects (see Section 4.1)

Distribution	Method	XMSE	XPE <sub>i</sub>			YMSE	YPE <sub>i</sub>		
			25th	50th	75th		25th	50th	75th
Gaussian	LGP	0.1242	0.1529	0.2847	0.7636	0.0076	0.0101	0.0205	0.0433
	GLMM-L	0.4182	0.3405	0.5843	1.283	0.0265	0.0278	0.0369	0.0577
	GLMM-Q	0.4323	0.3479	0.5990	1.319	0.0271	0.0285	0.0377	0.0584
	GEE-L	0.4168				0.0264			
	GEE-Q	0.4308				0.0272			
Non-Gaussian (mixture)	LGP	0.1272	0.1664	0.3166	0.9556	0.0078	0.0109	0.0228	0.0459
	GLMM-L	0.4209	0.3309	0.5943	1.364	0.0266	0.0280	0.0372	0.0589
	GLMM-Q	0.4373	0.3385	0.6118	1.404	0.0274	0.0287	0.0380	0.0597
	GEE-L	0.4227				0.0268			
	GEE-Q	0.4396				0.0277			

Simulation is based on 200 Monte Carlo runs with  $n = 100$  subjects per sample, generated from both Gaussian and non-Gaussian latent processes. Simulation is also repeated 1000 times for each of the parameters of interest. The mean and variance of the mean function  $\mu(t)$  and the variance  $\sigma^2(t)$  are given in Table 1. The mean and variance of the mean function  $\mu(t)$  and the variance  $\sigma^2(t)$  are given in Table 1. The mean and variance of the mean function  $\mu(t)$  and the variance  $\sigma^2(t)$  are given in Table 1. The mean and variance of the mean function  $\mu(t)$  and the variance  $\sigma^2(t)$  are given in Table 1.

and (30), evaluate the estimation error for the single eigenfunction in the model (noting that  $\int_{\mathcal{I}} \phi_1^2(t) dt = 1$ ),

$$EMSE = \int_{\mathcal{I}} \{\phi_1(t) - \hat{\phi}_1(t)\}^2 dt. \tag{31}$$

Using the same simulation design as in Section 4.1 and generating latent processes  $X(t; \delta) = \mu(t) + \delta \xi_1 \phi_1(t)$  for a range of  $\delta$ , estimate the 200 Gaussian and 200 non-Gaussian samples (as described before) for each of  $\delta = 0.5, 0.8, 1, 2$ . The Monte Carlo estimates of the mean and variance of  $\delta$  are presented in Table 2.

**Tab e 2.** Simulation results for the effect of the variation parameter  $\delta$

Distribution	$\delta$	XMSE	EMSE	XPE <sub>i</sub>			YMSE	YPE <sub>i</sub>		
				25th	50th	75th		25th	50th	75th
Normal	0.5	0.1106	0.7662	0.1188	0.1815	0.3366	0.0068	0.0077	0.0119	0.0205
	0.8	0.1205	0.3801	0.1430	0.2437	0.5710	0.0076	0.0094	0.0171	0.0338
	1	0.1280	0.2434	0.1513	0.2809	0.7857	0.0077	0.0101	0.0203	0.0431
	2	0.1616	0.0429	0.2025	0.3851	0.8137	0.0102	0.0144	0.0362	0.0752
Mixture	0.5	0.1134	0.7198	0.1243	0.1913	0.3651	0.0071	0.0081	0.0126	0.0217
	0.8	0.1258	0.3910	0.1498	0.2563	0.6691	0.0078	0.0100	0.0188	0.0366
	1	0.1323	0.2256	0.1624	0.2986	0.7944	0.0081	0.0113	0.0227	0.0450
	2	0.1633	0.0397	0.2041	0.3840	0.8140	0.0103	0.0158	0.0387	0.0768

Design and properties of the simulation are the same as in Table 1. EMSE denotes the average integrated mean squared error for estimating the single eigenfunction.

We find  $b$  an initial  $\delta$  of the error EMSE in estimating the eigenfunction on the interval  $[\delta, 1]$ . This is caused by the fact that  $\delta$  is small, increasing the accuracy in the observed data and the error of the panel of the underlying LGP, and the error becomes increasing difficult to estimate the eigenfunction. This is also observed in ordinary FPC analysis the error in estimating an eigenfunction is related to the error of the associated eigenvalue. The error of the eigenfunction can be estimated. Although the interval of  $\delta$  increases the error in predicting individual  $\mu_j(t)$ , this is in the prediction: for the prediction process  $X$ , this is because the accuracy of individual  $\mu_j(t)$  increases, the error of the binomial of the  $\mu_j(t)$  imposed constraints on the choice of this accuracy is affected in the prediction; for the  $\mu_j(t)$  process, the error increases in the choice, which is because the bias in the approximation has been reduced for the prediction and increasing  $\delta$ .



The error in estimating the mean function remains fairly stable as long as  $\delta \leq 1$ . This is especially true and not necessarily observed for the mean of prediction process  $X$ , since this mean estimates are not affected by an approximation error. We conclude that, namely  $\delta$  is large, it is actually has a small effect on the error in mean function estimates and a moderate effect on the error in individual prediction, and therefore has the long effect on the error in eigenfunction estimates does not play a role in the prediction for individual  $\mu_j(t)$ . The mean function estimates, as the effect is mitigated by the multiplication in  $\delta$ .

5. A case

Painful bilirubinemia (Morgan et al., 1994) is a rare but fatal chronic liver disease of unknown cause, with a prevalence of about 50 cases per million population. The data were collected between January 1974 and March 1984 by the Mayo Clinic (see also Appendix D of Fleming and Harrington (1991)). The patients were checked for the measurement of blood cholesteryl ester, a 6-month interval and annual health care diagnosis. However, since many individuals missed some of the checkups, the data are sparse and irregular. The number of measurements per subject and also the measurement time  $T_{ij}$  are individual.

To demonstrate the effectiveness of the methodology, we analyze the patient's history, who had a total of 10 years (3650 days) since he entered the study and eventually had not had a planned end of the 10 years. We carry out an analysis on the domain from 0 to 10 years, plotting the dynamic behavior of the presence of hepatomegaly (0, no; 1, yes), which is a longitudinal measurement. Benolli et al. (1984) and irregular measurements. The presence or absence of hepatomegaly is recorded on the data. The patient's age is between 42 years for the total of 429 binomially distributed observations, with a median of 11 measurements per subject.


We employ a logistic link function, and the model estimates of the mean and covariance function for the underlying process  $X(t)$  are displayed in Fig. 1. The mean function of the underlying process shows an increasing trend in the last 3000 days, especially after the beginning, and a decrease towards the end of the range of the data. We also provide pointwise bootstrap confidence intervals, which broaden (not necessarily) near the endpoints of the domain. The estimated covariance surface of  $X(t)$  displays rapidly decreasing correlation as the difference between measurements increases. With a variance function  $v(\mu) = \mu(1 - \mu)$ , the interval procedure for selecting the number of eigenfunctions and the variance parameter  $\gamma$  has been described in Section 3.2 yielded the choice  $M = 3$  for the number of components included and  $\gamma^2 = 1.91$  for the optimization parameter. The least one point



which are defined by equation (22), for the hepatic infection  $i(t)$



which are obtained by equation (23) for nine randomly selected values  $a, b, c, d, e, f, g, h, i$  shown in Figure 4. The predicted age-specific  $Y$  describes the time evolution of the probability of the presence of hepatitis for each individual; it is of increasing, decreasing, or mild oscillating decline.




The large projection in the direction of the respective eigenvectors shown in Figure 3(b).



**Fig. 2.** (a) Plot of  $PE(\gamma^2)$  values (26) of the final iteration versus corresponding candidate values of  $\gamma^2$ , where  $\hat{\gamma}^2$  minimizes  $PE(\gamma^2)$  and (b) FIC scores (28) for final iteration based on quasi-likelihood by using the binomial variance function for 10 possible leading eigenfunctions, where  $M = 3$  is the minimizing value (for the primary biliary cirrhosis data)

We find that the observed end of the predicted age-specific  $Y_i(t)$  agree well with the observed longitudinal binomial counts, and leave-one-out analysis using equation (24) confirmed this. In making the comparison between observed data and fitted probabilities, we need to keep in mind that the Bernoulli observations consist of 0's or 1's, whereas the fitted probabilities and dependent probabilities are constrained to be strictly between 0 and 1. Therefore, long-run analysis is expected for





(f)

(e)


(d)

(i)

(h)

(g)

process is not much influenced by age and  
age, but when income is non-linear, for age  
the conditional dependence of the effect  
the effect shape of the age income coefficient  
mean has older age and is associated





6. Discussion

The asymptotic theory for small  $\delta$  implies that the distribution in the latent process  $X$  is asymptotically normal, according to the asymptotic theory  $X(t) = \mu(t) + \delta Z(t)$ . We note that the small  $\delta$  asymptotic theory does not affect the methodology proposed, for which the value of  $\delta$  is not needed and plays no role. The estimation is proposed at a fixed age and age configuration for the initial LGP  $\tilde{X}$ , which is characterized by the mean function  $\nu(t)$  and covariance function  $\tau(s, t)$ , as defined in the previous section (8). However, bias estimates may be accounted for the estimation process, estimation and special prediction individual exposure adjustment for the case of large  $\delta$ .

$$U_{qr}(s, t) = \sum_{i:m_i \geq 2} \sum_{j,k:j \neq k} T_{ij}^q T_{ik}^r K_{ij}(s) K_{ik}(t),$$

$$\tilde{T}_{qr} = U_{qr} / U_{00},$$

$$\tilde{Z} = U_{00}^{-1} \sum_{i:m_i \geq 2} \sum_{j,k:j \neq k} Z_{ijk} K_{ij}(s) K_{ik}(t),$$

$$R = R_{20} R_{02} - R_{11}^2,$$

$Z_{ijk} = Y_{ij} Y_{ik}$ ,  $K_{ij}(t) = K\{(t - T_{ij})/h\}$ ,  $K$  is a kernel function and  $h$  a bandwidth. Of course, the eigenvalues of the same bandwidth operator  $\alpha$  and  $\beta$ ; the respective operators are band width for  $\tilde{\beta}$  to be large than for  $\alpha$ .

Both  $\alpha$  and  $\beta$  are convolutional, each has diagonal elements and convolutional elements. The data is in the  $i$ th block, i.e.  $\mathcal{B}_i = \{Y_{ij} \text{ for } 1 \leq i \leq m_i\}$ , are independent of one another, but the  $n$  blocks  $\mathcal{B}_1, \dots, \mathcal{B}_n$  are independent. Therefore, a least squares estimator of convolutional function (Rice and Sillerman, 1991) can be used to estimate the band width for each element.

### A and B: Periodic functions

Since the eigenvalues  $\tau(s, t)$  is symmetric, we have

$$\tau(s, t) = \sum_{j=1}^{\infty} \theta_j \psi_j(s) \psi_j(t), \tag{34}$$

where  $(\theta_j, \psi_j)$  are (eigenvalue, eigenfunction) pairs of a linear operator  $A$  in  $L^2$  which maps a function  $f$  to the function  $A(f)$ , which is defined by  $A(f)(s) = \int_{\mathcal{I}} \tau(s, t) f(t) dt$ . It is explained after equation (16) how the eigenvalues are obtained. Assuming that there is a finite number of the  $\theta_j$  are non-zero, the operator  $A$  will be positive semidefinite operator, eigenvalues  $\tau$  will be a positive covariance function, if and only if each  $\theta_j \geq 0$ . To ensure the operator is compact a condition (34) numerically and drop those elements that correspond to negative  $\theta_j$ , giving the eigenvalues

$$\tilde{\tau}(s, t) = \sum_{j \geq 1: \theta_j > 0} \theta_j \psi_j(s) \psi_j(t). \tag{35}$$

The modified eigenvalues  $\tilde{\tau}$  is not identical to  $\tau$  if one or more of the eigenvalues  $\theta_j$  are negative. In which case, the eigenvalues  $\tilde{\tau}$  has a  $L_2$ -accuracy than  $\tau$ , hence it is an eigenvalue of  $\tau$ .

*Theorem 1.* Under regularity conditions, it holds that

$$\int_{\mathcal{I}^2} (\tilde{\tau} - \tau)^2 \leq \int_{\mathcal{I}^2} (\tau - \tau)^2. \tag{36}$$

To prove this, let us show that condition (36) holds if and only if there are  $\tilde{\tau}$  is a non-trivial modification of  $\tau$ , i.e. when  $\tilde{\tau} \neq \tau$ . In the case where the high-frequency part of equation (34) is small, the low-frequency part of the eigenvalues has a high correlation with non-zero  $\theta_j$  are listed first, for  $1 \leq j \leq J$ , and  $\theta_j = 0$  only for  $j \geq J + 1$ . The sequence  $\psi_1, \dots, \psi_J$  is necessary and sufficient for the orthonormal, and the choice  $\psi_{J+1}, \psi_{J+2}, \dots$  of the orthonormal sequence  $\psi_1, \psi_2, \dots$  is orthonormal and also complete in the class of square integrable functions on  $\mathcal{I}$ .

We may therefore express the covariance  $\tau$  in terms of the eigenvalues, as a convolutional expansion in a generalized Fourier series:

$$\tau(s, t) = \sum_{j=1}^{\infty} \sum_{k=1}^{\infty} a_{jk} \psi_j(s) \psi_k(t), \tag{37}$$

where  $a_{jk} = \int_{\mathcal{I}^2} \tau(s, t) \psi_j(s) \psi_k(t) ds dt$ . Equation (34), (35) and (37) imply that

$$\int_{\mathcal{I}^2} (\tilde{\tau} - \tau)^2 = \sum_{j,k:j \neq k} a_{jk}^2 + \sum_{j=1}^{\infty} (a_{jj} - \tilde{\theta}_j)^2,$$

$$\int_{\mathcal{I}^2} (\tau - \tau)^2 = \sum_{j,k:j \neq k} a_{jk}^2 + \sum_{j=1}^{\infty} (a_{jj} - \tilde{\theta}_j)^2,$$

$$\sigma_{ikl} \equiv \text{cov}(\tilde{X}_{ik}, \tilde{X}_{il}) = \sum_j \theta_j \psi_j(T_{ik}) \psi_j(T_{il}) + \delta_{kl} \frac{\gamma^2 v\{g\{\mu(T_{ik})\}\}}{g^{(1)}\{\mu(T_{ik})\}^2},$$

where  $\delta_{kl} = 1$  if  $k=l$  and 0 otherwise, and

$$d_i \equiv \tilde{X}_i - E(\tilde{X}_i) = \left( \frac{Y_{i1} - g\{\mu(T_{i1})\}}{g^{(1)}\{\mu(T_{i1})\}}, \dots, \frac{Y_{im_i} - g\{\mu(T_{im_i})\}}{g^{(1)}\{\mu(T_{im_i})\}} \right)^T.$$

Denote  $\text{cov}(\tilde{X}_i, \tilde{X}_i) = \Sigma_i = (\sigma_{ikl})_{1 \leq k, l \leq m_i}$ . Then the explicit form of the matrices  $A_{ij}$  in equation (21) is given by

$$E(\xi_{ij} | Y_{i1}, \dots, Y_{im_i}) = \theta_j \psi_{ij} \Sigma_i^{-1} d_i, \tag{39}$$

where  $\psi_{ij}$  is the  $j$ th component of the  $i$ th row of the matrix  $\Psi$  in equation (15),  $\gamma$  is the scale parameter in equation (27), and  $\theta_j$  and  $\psi_j$  are the coefficients of the  $j$ th component of the  $i$ th row of the matrix  $\Theta$  in equation (27).

**References**

Boente, G. and Faiman, R. (2000) Kernel-based functional principal components. *Statist. Probab. Lett.*, **48**, 335–345.

Cheng, M. Y., Hall, P. and Tien, D. M. (1997) On the linkage of local linear regression. *Statist. Comput.*, **7**, 11–17.

Chio, J.-M. and Müller, H.-G. (2005) Estimation of the regression function: empirical inference for clustered and longitudinal data. *J. R. Statist. Soc. B*, **67**, 531–553.

Chio, J. M., Müller, H. G. and Wang, J. L. (2004) Functional regression model. *Statist. Sin.*, **14**, 675–693.

Diggle, P. J., Tan, J. A. and Moeed, R. A. (1998) Model-based geostatistics (in Hindi). *Appl. Statist.*, **47**, 299–350.

Fan, J. (1993) Local linear regression and its minimax efficiency. *Ann. Statist.*, **21**, 196–216.

Fleming, T. R. and Harrington, D. P. (1991) *Counting Processes and Survival Analysis*. New York: Wiley.

Hajmiri, R., Jacomini-Gadda, H. and Commenges, D. (2003) A latent process model for joint modeling of event and marker. *Lifetime Data Anal.*, **9**, 331–343.

Heagerty, P. J. (1999) Marginalized logistic-normal model for longitudinal binary data. *Biometrics*, **55**, 688–698.

Heagerty, P. J. and Kurland, B. F. (2001) Marginalized multivariate likelihood estimation and generalized linear mixed model. *Biometrika*, **88**, 973–985.

Heagerty, P. J. and Zeger, S. L. (2000) Marginalized multilevel model and likelihood inference. *Statist. Sci.*, **15**, 1–26.

Jamshidi, G., Hajmiri, T. G. and Sgambati, C. A. (2001) Principal component model for sparse functional data. *Biometrika*, **87**, 587–602.

Jamshidi, G. and Sgambati, C. A. (2003) Classification of sparse functional data. *J. Am. Statist. Ass.*, **98**, 397–408.

Johndrow, V. and Sridhar, B. (2002) Analyzing longitudinal count data with hierarchical structure. *Biometrika*, **89**, 389–399.

Kippenhahn, M. and Heckman, N. (1989) A variational genetic model for growth, shape, reaction norm, and other infinite-dimensional characteristics. *J. Math. Biol.*, **27**, 429–450.

Munz, P. A., Dickson, E. R., Van Dam, G. M., Malinchoc, M., Gambich, P. M., Langford, A. L. and Gipson, C. H. (1994) Proinflammatory cytokines: prediction of hospital mortality based on repeated plasma levels. *Hepatology*, **20**, 126–134.

Poahmadi, M. (2000) Marginalized multivariate likelihood estimation of generalized linear model for multivariate normal covariance matrix. *Biometrika*, **87**, 425–435.

Poon, C., Jacomini-Gadda, H., Touloukian, J. M. G., Ganiyu, J. and Commenges, D. (2006) A nonlinear model for longitudinal cognitive decline in aging memory impairment and longitudinal data. *Biometrics*, **62**, 1014–1024.

Ramsey, J. and Silverman, B. (2002) *Applied Functional Data Analysis*. New York: Springer.

Ramsey, J. and Silverman, B. (2005) *Functional Data Analysis*, 2nd edn. New York: Springer.

Rice, J. (2004) Functional and longitudinal data analysis: prediction of mortality. *Statist. Sin.*, **14**, 631–647.

Rice, J. A. and Silverman, B. W. (1991) Estimation of the mean and covariance structure of nonparametrically estimated functional data. *J. R. Statist. Soc. B*, **53**, 233–243.

Rice, J. and Wang, C. (2000) Nonparametric mixed effect model for nested longitudinal data. *Biometrics*, **57**, 253–259.

Seifried, B. and Gasser, T. (1996) Functional data analysis of local polynomial regression: analysis and simulation. *J. Am. Statist. Ass.*, **91**, 267–275.

- Shi, M., Wei, R. E. and Talo, J. M. G. (1996) An analysis of paediatric CD4 counts for acquired immune deficiency syndrome using flexible random effects. *Appl. Statist.*, **45**, 151–163.
- Sani, Ali, J. G. and Lee, J. J. (1998) Nonparametric regression analysis of longitudinal data. *J. Am. Statist. Ass.*, **93**, 1403–1418.
- Yao, F., Mullen, H. G., Clifford, A. J., Deeks, S. R., Follett, J., Lin, Y., Buchholz, B. A. and Vogel, J. S. (2003) Shrinkage estimation for functional principal component coefficients in application to the population kinetics of plasma folate. *Biometrics*, **59**, 676–685.
- Yao, F., Mullen, H. G. and Wang, J. L. (2005) Functional data analysis for sparse longitudinal data. *J. Am. Statist. Ass.*, **100**, 577–590.
- Zhao, X., Ma, J., Jones, J. S. and Wells, M. T. (2004) The functional data analysis of longitudinal data. *Statist. Sin.*, **14**, 789–808.